

ON THE ASYMPTOTIC BEHAVIOUR OF THE ZEROS OF SOLUTIONS OF ONE FUNCTIONAL-DIFFERENTIAL EQUATION WITH RESCALING

GREGORY DERFEL, PETER J. GRABNER^{†*}, AND ROBERT F. TICHY^{‡*}

Dedicated to Heinz Langer on the occasion of his 80th birthday.

ABSTRACT. We study the asymptotic behaviour of the solutions of a functional- differential equation with rescaling, the so-called pantograph equation. From this we derive asymptotic information about the zeros of these solutions.

1. INTRODUCTION

1.1. Historical remarks. The importance of functional and functional-differential equations with rescaling is being increasingly recognised in the last decades, as is their relevance to a wide range of application areas. Such equations – functional and functional-differential, linear and nonlinear – appear as adequate tools in a number of phenomena that display some kind of self-similarity. It is next to impossible to describe all recent activities in this area. For general references and bibliography we refer the reader to survey papers by Derfel and co-authors [1, 6, 7].

One of the best known examples of equations with rescaling is the celebrated *pantograph equation*:

$$(1.1) \quad y'(z) = ay(\lambda z) + by(z).$$

This equation was introduced by Ockendon & Tayler [20] as a mathematical model of the overhead current collection system on an electric locomotive. (The term ‘pantograph equation’ was coined by Iserles [13].) This equation and its ramifications have emerged in a striking range of applications, including number theory [18], astrophysics [3], queues & risk theory [10], stochastic games [9], quantum theory [23], population dynamics [12], and graph theory [21]. The common feature of all such examples is some self-similarity of the system under study.

Date: December 20, 2016.

2000 Mathematics Subject Classification. Primary: ; Secondary:

Key words and phrases. functional differential equation, zeros.

[†] This author is supported by the Austrian Science Fund FWF project F5503 (part of the Special Research Program (SFB) “Quasi-Monte Carlo Methods: Theory and Applications”).

[‡] This author is supported by the Austrian Science Fund FWF project F5510 (part of the Special Research Program (SFB) “Quasi-Monte Carlo Methods: Theory and Applications”).

^{*} These authors are supported by the Doctoral Program “Discrete Mathematics” W1230.

In 1972, Morris, Feldstein and Bowen [19] studied functional-differential equations (FDE) of the form:

$$(1.2) \quad y'(z) = \sum_{k=1}^{\ell} a_k y(\lambda_k z)$$

with $1 > \lambda_{\ell} > \lambda_{\ell-1} > \dots > \lambda_1 > 0$, $a_1, \dots, a_{\ell} \in \mathbb{C}$. They were able to obtain deep results about the existence, uniqueness, and asymptotic behaviour of solutions of (1.2) in the complex plane \mathbb{C} . (For more about FDE in the complex plane see also [8].)

Among other things, Morris *et al.* gave a detailed analysis of (1.1) in the special case $a = -1, b = 0$, i.e.,

$$(1.3) \quad y'(z) = -y(\lambda z)$$

In particular, they proved that (1.1), supplemented by the initial condition

$$(1.4) \quad y(0) = 1$$

has the unique solution

$$(1.5) \quad y(z) = \sum_{n=0}^{\infty} (-1)^n \lambda^{\frac{n(n-1)}{2}} \frac{z^n}{n!}.$$

Moreover, $y(z)$ is an entire function of order zero, and it has infinitely many positive zeros, but no other zeros in the complex plane. The entire function (1.5) is sometimes called the *deformed exponential function* (see Sokal [22]).

A number of conjectures on the zeros $0 < t_0 < t_1 < t_2 \dots$ of (1.5) have been made by Morris, Feldstein and Bowen [19] and by Iserles [13]. In particular, Morris, Feldstein and Bowen conjectured that

$$(1.6) \quad \lim_{n \rightarrow \infty} t_{n+1}/t_n = 1/\lambda := q$$

Also, in what follows we shall use the notation $q := 1/\lambda > 1$.

It is notable that in 1973, independently of [19], Robinson, in his paper on counting of acyclic digraphs [21], derived the same FDE as (1.3) and conjectured

$$(1.7) \quad t_n = (n+1)q^n + o(q^n),$$

Clearly, (1.7) is stronger than (1.6). In 2000, Langley [15] resolved Morris's *et al.* conjecture by proving that

$$(1.8) \quad t_n = nq^{n-1}(\gamma + o(1)),$$

where γ is a positive constant. In 2005, Grabner and Steinsky [11], independently of [15], proved a weaker form of Robinson's conjecture: namely, there exists k_0 such that

$$(1.9) \quad t_{k_0+k} = (k+1)q^k + o(q^k/k^{(1-\varepsilon)}),$$

for all $\varepsilon > 0$.

Recently, Zhang [24] proved that $\gamma = 1$ in (1.8) and, moreover,

$$(1.10) \quad t_n = nq^{n-1}(1 + \psi(q)n^{-2} + o(n^{-2})),$$

where $\psi(q)$ is the generating function of the sum-of-divisors function $\sigma(k)$ [15]. Also, he derived an asymptotic formula for the oscillation amplitude A_n of $y(x)$, i.e., $A_n = |y(qt_n)|$.

1.2. Main results. All aforementioned results were concerned with the analytic function (1.5), which is the unique solution of the Cauchy problem (1.3)–(1.4). In contrast to this, in the present paper we deal with *all solutions* of (1.3), not necessarily also satisfying (1.4), but rather *defined on an arbitrary half-line*.

The following natural definition is commonly accepted in the theory of functional-differential equations:

Definition 1.1. If x_0 is a real number, then a real or complex function $y(x)$, defined and continuous for $x \geq \lambda x_0$, is said to be a solution of (1.3) for $x \geq x_0$, if it satisfies (1.3) for all $x \geq x_0$.

Thus, instead of the Cauchy problem for ODE, we have an *initial value* problem for FDE of *retarded type* (i.e., $0 < \lambda < 1$): for an arbitrary *initial function* φ defined on $[\lambda x_0, x_0]$, a solution of the initial value problem is a function $y(x)$ defined and continuous for $x \geq \lambda x_0$, which satisfies (1.3) and the initial condition:

$$(1.11) \quad y(x) = \varphi(x), \quad x \in [\lambda x_0, x_0].$$

Thus, the general solution of FDEs normally consists of an *infinite family of solutions*, depending on an *arbitrary function*.

Evidently, any function that is continuous in $[\lambda x_0, x_0]$ can be continued uniquely to a solution for $x \geq x_0$, and it makes sense to discuss the asymptotic behaviour of all solutions as $x \rightarrow \infty$.

The aim of this paper is to show that the above stated asymptotic behaviour of zeros is true not only for the analytic solution (1.5), but for *all* solutions of (1.3).

Namely, in Section 3, below, we prove that for every solution $y(x)$ of (1.3), the following asymptotic formula for zeros t_n is valid:

$$(1.12) \quad t_n = nq^{n-1} \left(\gamma + \mathcal{O}\left(\frac{\log k}{k}\right) \right)$$

It is worth noting that all solutions of (1.3), except for (1.5), are *non-analytic* ones at $z = 0$, but (1.12) remains true for all of them.

In summary, we can say that the asymptotic behaviour (1.12) is an *intrinsic property of the equation* (1.3) itself, and only the constant γ in (1.12) depends on the specific solution.

Our second objective (see Section 2) is an asymptotic analysis of the solutions of (1.3).

In Section 2.1 we derive an asymptotic formula of de Bruijn, Kato and McLeod type by a method different from [2] and [14].

Furthermore, in Section 2.2 we give a brief summary of related results from [4] and [5] for *general FDE of higher order with several scaling factors*

$$(1.13) \quad y^{(m)}(x) = \sum_{j=0}^{\ell} \sum_{k=0}^{m-1} a_{jk} y^{(k)}(\alpha_j t + \beta_j),$$

where all α_j are less than 1 in modulus i.e. $|\alpha_j| < 1$.

2. ASYMPTOTIC BEHAVIOUR OF ANALYTIC SOLUTIONS

2.1. Asymptotic behaviour of the solution of FDE $y'(z) = -y(\lambda z)$. We first observe that the equation

$$(2.1) \quad y'(z) = ay(\lambda z)$$

can be simplified to

$$(2.2) \quad g'(z) = -g(\lambda z);$$

i.e. every solution of (2.1) can be written as $y(z) = g(-az)$ for a solution g of (2.2).

We start with an ansatz as a power series

$$g(z) = \sum_{n=0}^{\infty} g_n z^n,$$

which gives the recursion

$$(n+1)g_{n+1} = -\lambda^n g_n.$$

From this we obtain

$$(2.3) \quad g(z) = \sum_{n=0}^{\infty} (-1)^n \lambda^{\binom{n}{2}} \frac{z^n}{n!},$$

if we assume $g_0 = 1$.

In order to study the asymptotic behaviour of $g(z)$ for large z , we transform (2.3) into an integral representation inspired by the inversion formula for the Mellin transform

$$(2.4) \quad g(z) = \frac{1}{2\pi i} \int_{\mathcal{H}} \Gamma(s) \lambda^{\binom{-s}{2}} z^{-s} ds = \frac{1}{2\pi i} \int_{\mathcal{H}} \frac{\pi}{\sin(\pi s)} \frac{\lambda^{\binom{-s}{2}} z^{-s}}{\Gamma(1-s)} ds,$$


$$(2.5) \quad g(-z) = \frac{1}{2\pi i} \int_{\mathcal{H}} \pi \cot(\pi s) \frac{\lambda^{\binom{-s}{2}} z^{-s}}{\Gamma(1-s)} ds,$$

where \mathcal{H} is a contour encircling the negative real axis counterclockwise, like the Hankel-contour used in the theory of the Γ -function (see Figure 1). We will use the representation (2.4) for deriving an asymptotic formula for $g(z)$ for $|\arg(z)| \leq \pi - \varepsilon$ (for $\varepsilon > 0$), whereas (2.5) will be used for the asymptotic of $g(z)$ for $|\arg(-z)| \leq \pi - \varepsilon$. These representations can be proved by residue calculus and taking care of the growth order of the integrand along the contour. In the case $\lambda = \frac{1}{2}$ a similar representation was used in [11].

Similarly, the representation

$$g(z) = \frac{1}{2\pi i} \left(\int_{-\infty-i}^{\infty-i} - \int_{-\infty+i}^{\infty+i} \right) \Gamma(s) \lambda^{\binom{-s}{2}} z^{-s} ds$$

can be shown. Here the contour of integration is deformed into two horizontal lines above and below the real axis.

FIGURE 1. The contour of integration \mathcal{H}

Theorem 2.1. *Let g be the entire solution of (2.2) with $g(0) = 1$. Then for $\varepsilon > 0$ and $|\arg(z)| \leq \pi - \varepsilon$ and $z \rightarrow \infty$ the asymptotic expansion*

$$(2.6) \quad g(z) \sim C z^A \log(z)^B \exp \left(-\frac{1}{2 \log \lambda} (\log(z) - \log(\log(z)))^2 \right) \\ \times H \left(\frac{1}{\log \lambda} (\log(z) - \log(\log(z))) - \frac{1}{2} + \frac{\log(-\log \lambda)}{\log \lambda} \right)$$

holds, where the periodic function H of period 2 is given by the Fourier series

$$(2.7) \quad H(x) = \sqrt{\frac{2\pi}{-\log \lambda}} \sum_{k \in \mathbb{Z}} e^{\frac{(2k+1)^2 \pi^2}{2 \log \lambda}} e^{\pi i (2k+1)x},$$

and

$$(2.8) \quad A = \frac{1}{2} - \frac{1}{\log \lambda} - \frac{\log(-\log \lambda)}{\log \lambda} \\ B = \frac{\log(-\log \lambda)}{\log \lambda} - 1 \\ C = \exp \left(\frac{1}{2} - \frac{\log \lambda}{8} + \log(-\log \lambda) - \frac{\log(-\log \lambda)}{\log \lambda} - \frac{(\log(-\log \lambda))^2}{2 \log \lambda} - \frac{1}{2} \log(2\pi) \right).$$

Remark 2.1. This theorem extends the real asymptotic of **all** solutions on \mathbb{R}^+ given in [14] to the complex asymptotic of the **entire** solution in an angular region avoiding the negative real axis.

Theorem 2.2. *Let g be the entire solution of (2.2) with $g(0) = 1$. Then for $\varepsilon > 0$ and $|\arg(-z)| \leq \pi - \varepsilon$ and $z \rightarrow \infty$ the asymptotic expansion*

$$(2.9) \quad g(-z) \sim C z^A \log(-z)^B \exp \left(-\frac{1}{2 \log \lambda} (\log(-z) - \log(\log(-z)))^2 \right) \\ \times K \left(\frac{1}{\log \lambda} (\log(-z) - \log(\log(-z))) - \frac{1}{2} + \frac{\log(-\log \lambda)}{\log \lambda} \right)$$

holds, where the periodic function K of period 1 is given by the Fourier series

$$(2.10) \quad K(x) = \sqrt{\frac{2\pi}{-\log \lambda}} \sum_{k \in \mathbb{Z}} e^{\frac{2k^2 \pi^2}{\log \lambda}} e^{2\pi i k x},$$

and A , B , and C are given by (2.8).

Proof of Theorem 2.1. We apply a saddle point approximation combined with residue calculus to the second integral representation given in (2.4). For this purpose, we consider

$$(2.11) \quad \frac{\lambda^{\binom{-s}{2}} z^{-s}}{\Gamma(1-s)} = \exp \left(\frac{s(s+1)}{2} \log \lambda - s \log(z) - \log(-s) - \log(\Gamma(-s)) \right).$$

We find the saddle point as the stationary point of the argument of the exponential:

$$(2.12) \quad \left(s + \frac{1}{2} \right) \log \lambda - x - \frac{1}{s} + \psi(-s) = 0,$$

where $\psi = \frac{\Gamma'}{\Gamma}$; for simplicity, we set $x = \log(z)$. From [17] we infer

$$\psi(-s) = \log(-s) + \frac{1}{2s} + \mathcal{O}\left(\frac{1}{s^2}\right),$$

valid for $|\arg(-s)| \leq \pi - \varepsilon$ for $\varepsilon > 0$. This shows that (2.12) has a unique solution $\sigma(x)$ satisfying

$$(2.13) \quad \sigma(x) = \frac{1}{\log \lambda} (x - \log(x)) - \frac{1}{2} + \frac{\log(-\log \lambda)}{\log \lambda} + \mathcal{O}(\log(x)/x),$$

which lies close to the negative real axis; notice that the imaginary part of $\sigma(x)$ is bounded by $\frac{\pi}{|\log \lambda|}$. Around $s = \sigma(x)$ we have the following approximation

$$(2.14) \quad \frac{s(s+1)}{2} \log \lambda - sx - \log(s) - \log(\Gamma(-s)) \\ = \frac{\sigma(x)(\sigma(x)+1)}{2} \log \lambda - x\sigma(x) - \log(-\sigma(x)) - \log(\Gamma(-\sigma(x))) \\ + \frac{1}{2} \left(\log \lambda + \frac{1}{\sigma(x)^2} - \psi'(-\sigma(x)) \right) (s - \sigma(x))^2 + \mathcal{O}(x^{-2}(s - \sigma(x))^3)$$

for $s - \sigma(x) = \mathcal{O}(x^\alpha)$ for $\alpha < \frac{2}{3}$.

Inserting the approximation (2.14) into the integral representation (2.4) (and splitting the range of integration into $|s - \sigma(x)| \leq x^\alpha$ and $|s - \sigma(x)| \geq x^\alpha$) yields

$$(2.15) \quad g(z) = \frac{\lambda^{\binom{-\sigma(x)}{2}}}{\Gamma(1 - \sigma(x))} e^{-x\sigma(x)} \frac{1}{2\pi i} \oint_{R_x} \exp\left(\frac{1}{2}(s - \sigma(x))^2 \log \lambda\right) \frac{\pi}{\sin(\pi s)} ds (1 + \mathcal{O}(x^{2\alpha-1})) \\ + \frac{1}{2\pi i} \int_{R'_x} \Gamma(s) \lambda^{\binom{-s}{2}} z^{-s} ds,$$

if we choose $\alpha < \frac{1}{2}$. Here R_x denotes the positively oriented rectangle with corners $\sigma(x) \pm |x|^\alpha \pm \frac{2\pi i}{\log \lambda}$ and R'_x denotes the remaining part of the dashed contour in Figure 1.

The first integral in (2.15) can be evaluated by residue calculus

$$(2.16) \quad \frac{1}{2\pi i} \oint_{R_x} \exp\left(\frac{1}{2}(s - \sigma(x))^2 \log \lambda\right) \frac{\pi}{\sin(\pi s)} ds \\ = \sum_{\substack{n \in \mathbb{Z} \\ |n - \sigma(x)| < |x|^\alpha}} (-1)^n e^{\frac{1}{2}(n - \sigma(x))^2 \log \lambda} = \sum_{n \in \mathbb{Z}} (-1)^n e^{\frac{1}{2}(n - \sigma(x))^2 \log \lambda} + \mathcal{O}\left(e^{\frac{1}{2}|x|^{2\alpha} \log \lambda}\right).$$

This series represents a continuous periodic function of period 2, whose Fourier coefficients can be computed by a variant of Poisson's summation formula

$$(2.17) \quad H(y) = \sum_{n \in \mathbb{Z}} (-1)^n e^{\frac{1}{2}(n - y)^2 \log \lambda} = \frac{\sqrt{2\pi}}{\sqrt{-\log \lambda}} \sum_{k \in \mathbb{Z}} e^{\frac{(2k+1)^2 \pi^2}{2 \log \lambda}} e^{i\pi(2k+1)y}.$$

For estimating the remaining integral over R'_x in (2.15), we use the estimates (cf. [17])

$$|\Gamma(t \pm iC)| \leq \sqrt{2\pi} (t^2 + C^2)^{\frac{1}{2}(t - \frac{1}{2})} e^{-\frac{\pi C}{2} + \frac{1}{6C}} \\ |\Gamma(-t \pm iC)| \leq \frac{\pi}{|C| \cosh(\pi C)^{\frac{1}{2}} \Gamma(t)}$$

valid for $t \geq 1$ and $C > 0$.

Inserting the asymptotic information about $\sigma(x)$ from (2.13) into (2.11) and using Stirling's formula for the Γ -function yields

$$\frac{\lambda^{\binom{-\sigma(x)}{2}} e^{-x\sigma(x)}}{\Gamma(1 - \sigma(x))} = C e^{Ax} x^B \exp\left(-\frac{1}{2 \log \lambda} (x - \log(x))^2\right),$$

with A , B , and C as given in (2.8). □

Proof of Theorem 2.2. For the proof of (2.9) we use the integral representation (2.5) and argue along the same lines as in the proof of Theorem 2.1. The only technical difference is that the residues of $\pi \cot(\pi s)$ are all equal to 1, which avoids the sign change occurring in (2.16). □

2.2. FDE of higher order with compressed arguments. In this section we consider the general FDE with rescaling

$$(2.18) \quad y^{(m)}(x) = \sum_{j=0}^l \sum_{k=0}^{m-1} a_{jk} y^{(k)}(\alpha_j t + \beta_j),$$

where $a_{jk} \in \mathbb{C}$ and $\alpha_j, \beta_j \in \mathbb{R}$. From here on we consider, solutions of (2.18) defined on the whole real line \mathbb{R} , when there exists at least one $\beta_j \neq 0$, and possibly defined on a half-line \mathbb{R}_+ , or \mathbb{R}_- , when all $\beta_j = 0$.

For such equations, it is hard to expect the existence of an asymptotic formula similar to (2.6). However, we can derive sharp estimates from above and below for solutions of (2.18).

Below, we give a brief summary of the related results from [4] and [5].

Denote:

$$(2.19) \quad \alpha = \min_{0 \leq j \leq l} |\alpha_j|, \quad A = \max_{0 \leq j \leq l} |\alpha_j|.$$

Then:

- (i) Every solution $y(x)$ of (2.18) is an analytic function, that can be extended as an entire function $y(z)$ of order zero in \mathbb{C} .
- (ii) Every solution of (2.18) satisfies the estimate

$$(2.20) \quad |y(z)| \leq C \exp\{\gamma \log^2(1 + |z|)\}, \quad z \in \mathbb{C},$$

for some $C > 0$, and

$$(2.21) \quad \gamma > m/(2|\log A|)$$

- (iii) Every solution of (2.18) is unbounded *on any ray* emanating from the origin $z = 0$ (as it is an entire function of order zero). In particular, every solution is unbounded on \mathbb{R}_+ and \mathbb{R}_- .

This result cannot be strengthened in general due to the existence of polynomial solutions. It was proved in [4] and [5] that:

- (iv) A necessary and sufficient condition for the existence of polynomial solutions of (2.18) is that:

$$(2.22) \quad \sum_{j=0}^l a_{j0} \alpha_j^n = 0$$

for some $n \in \mathbb{N}$.

Under the assumption that (2.18) has no polynomial solutions, and $\beta_j = 0$ for all j , one can prove a result stronger than (iii). Roughly speaking, every nontrivial solution of (2.18) grows as $|z| \rightarrow \infty$ faster than $\exp\{\gamma \log^2 |z|\}$ for some $\gamma > 0$. More precisely:

- (v) Every solution of (2.18), which *at least on one ray* emanating from the origin satisfies (2.20) for some $C > 0$, and

$$(2.23) \quad \gamma < 1/(2|\log \alpha|)$$

vanishes identically.

Remark 2.2. The above stated results, formulated for the equation (2.18), with *constant coefficients*, remain true with minor changes also for FDE of higher order, with several compressed arguments and *polynomial coefficients*:

$$(2.24) \quad y^{(m)}(x) = \sum_{j=0}^l \sum_{k=0}^{m-1} \sum_{n=0}^r a_{jkn} x^\nu y^{(k)}(\alpha_j t + \beta_j),$$

3. ASYMPTOTIC BEHAVIOUR OF THE ZEROS

The central result of this section is the following theorem.

Theorem 3.1. *Let $y(x)$ be a solution of (1.3), and $0 < x_0 < x_1 < x_2 < \dots$ be the zeros of $y(x)$. Then there exists a positive constant γ , such that:*

$$(3.1) \quad x_n = nq^{n-1} \left(\gamma + \mathcal{O}\left(\frac{\log k}{k}\right) \right).$$

The proof is based on a result of Kato & McLeod [14] and de Bruijn [2] on the asymptotic behaviour of *all solutions* of the equation

$$(3.2) \quad y'(z) = ay(\lambda z)$$

and Lemma 3.1 below.

According to Kato and McLeod [14, Theorem 7(iii)] and de Bruijn [2, Sections 1.3-1.4] every solution of (3.2) has the following asymptotic behaviour

$$(3.3) \quad y(x) = x^{A_1} \log(x)^{B_1} \exp \left(-\frac{1}{2 \log \lambda} (\log(x) - \log(\log(x)))^2 \right) \\ \times \left(h \left(\frac{1}{\log \lambda} (\log(x) - \log(\log(x))) \right) + o(1) \right),$$

where

$$(3.4) \quad A_1 = \frac{1}{2} - \frac{1}{\log \lambda} - \frac{\log(-a \log \lambda)}{\log \lambda}, \\ B_1 = \frac{\log(-a \log \lambda)}{\log \lambda} - 1,$$

and $h(x)$ is a periodic function of period $\log q = |\log \lambda|$, with some additional assumptions on its Fourier coefficients (see [14, (6.2)–(6.3)]).

Remark 3.1. Notice that this asymptotic behaviour is in accordance with the behaviour stated in Theorems 2.6 and 2.2. If the value $-a \log \lambda$ is negative, the complex values of the power $(x/\log(x))^{-\log(-a \log \lambda)/\log \lambda}$ are compensated by complex values of the periodic

function $h(x)$ and a doubling of the period to accommodate the sign change. This is the explanation of the fact that the periodic function in Theorem 2.6 has period 2.

Lemma 3.1. *Let G be a periodic function of period 2, x_0 the minimal positive zero of G , and all zeros of G located at the points: $x_0 + k$; $k = 0, 1, 2, \dots$. Then the zeros of the function $F(x) := G\left(\frac{1}{\log \lambda}(\log(x) - \log(\log(x)))\right)$ have the following asymptotic behaviour*

$$(3.5) \quad x_n = nq^{n-1} \left(\gamma + \mathcal{O}\left(\frac{\log k}{k}\right) \right)$$

Proof of Lemma 3.1. Let us observe first that the zeros of $F(x)$ are located at the points x , such that

$$(3.6) \quad \log x - \log x \log x = (x_0 + k) \log q$$

We shall seek solutions of (3.6) of the form

$$(3.7) \quad x_k = x(k) = C(k)kq^k,$$

where $C(k)$ is an unknown function of no more than power growth, i.e., there exists $\alpha > 0$ such that $C(k) = \mathcal{O}(k^\alpha)$. To prove Lemma 3.1 it is enough to show that

$$(3.8) \quad C(k) = \gamma + \mathcal{O}\left(\frac{\log k}{k}\right),$$

where γ is positive constant. For that, substitute (3.7) in (3.6). It follows from (3.7) that

$$(3.9) \quad \log x = k \log q \left(1 + \frac{\log k}{k \log q} + \frac{\log C(k)}{k \log q} \right),$$

and

$$(3.10) \quad \log \log x = \log k + \frac{\log k}{k \log q} + \frac{\log C(k)}{k \log q} + \log \log q + o\left(\frac{\log k}{k \log q}\right),$$

Combining (3.6), (3.9) and (3.10), after some elementary calculations we obtain that

$$(3.11) \quad \left(1 - \frac{1}{k \log q} \right) \log C(k) = x_0 \log q + \log \log q + \frac{\log k}{k \log q} + o\left(\frac{\log k}{k \log q}\right)$$

Next, let $k \rightarrow \infty$, then from (3.11) we obtain

$$(3.12) \quad \log C(k) = (x_0 \log q + \log \log q) + \mathcal{O}\left(\frac{\log k}{k}\right),$$

or

$$(3.13) \quad C(k) = q^{x_0} \log q \left(1 + \mathcal{O}\left(\frac{\log k}{k}\right) \right).$$

Finally, (3.13) implies (3.8) and (3.5). □

Proof of Theorem 3.1. First, we apply the asymptotic formula (3.3) to the solutions of (1.3). Observe that in this case $a = -1$ and $\log \lambda = -\log q < 0$, and therefore the expression $\log(-a \log \lambda) = \log(\log \lambda)$ in (3.4) is a complex number.

Having this in mind, we can rewrite (3.3) and (3.4) in the form:

$$(3.14) \quad y(x) = x^{A_2} \log(x)^{B_2} \exp \left(\frac{1}{2 \log q} (\log(x) - \log(\log(x)))^2 \right) \\ \times \left(\cos \frac{\pi}{\log q} (\log(x) - \log(\log(x))) + i \left(\sin \frac{\pi}{\log q} (\log(x) - \log(\log(x))) \right) \right) \\ \times \left(h \left(\frac{1}{\log q} (\log(x) - \log(\log(x))) \right) + o(1) \right),$$

where

$$(3.15) \quad A_2 = \frac{1}{2} + \frac{1}{\log q} + \frac{\log(\log q)}{\log q} \\ B_2 = -1 - \frac{\log(\log q)}{\log q},$$

Both the real part and the imaginary parts of (3.14) provide asymptotic formulas for solutions of (1.3), and the assumptions of Lemma 3.1 are fulfilled.

Then, in accordance with Lemma 3.1, there exists k_0 such that for all $k = 0, 1, 2, \dots$

$$(3.16) \quad x_{k_0+k} = kq^{k-1} \left(\gamma_1 + \mathcal{O}\left(\frac{\log k}{k}\right) \right),$$

Now (3.8) follows from (3.16), and from (3.8) finally we obtain (3.1). \square

4. THE CASE $b \neq 0$

The behaviour of the entire solution of (1.1) changes considerably, if $b \neq 0$. This is quite obvious from the observation that the equation can be viewed as a perturbed differential equation $y' = by$. Thus we expect an asymptotic behaviour of the form $f(x) \sim Ce^{bx}$ for $\Re(bx) \rightarrow +\infty$. We again start with an ansatz as a power series

$$f(z) = \sum_{n=0}^{\infty} f_n z^n,$$

from which we derive the recurrence formula

$$(n+1)f_{n+1} = (a\lambda^n + b)f_n \text{ for } n \geq 0.$$

This gives

$$(4.1) \quad f_n = \frac{b^n}{n!} \prod_{k=0}^{n-1} \left(1 + \frac{a}{b} \lambda^k \right),$$

if we set $f_0 = 1$. In order to simplify notation, we introduce

$$Q_\lambda(\alpha) = \prod_{k=0}^{\infty} (1 + \alpha \lambda^k).$$

This gives

$$(4.2) \quad f(z) = Q_\lambda \left(\frac{a}{b} \right) \sum_{n=0}^{\infty} \frac{(bz)^n}{n! Q_\lambda \left(\frac{a}{b} \lambda^n \right)},$$

if $\frac{a}{b} \lambda^n \neq -1$ for all $n \in \mathbb{N}$. In the case that $\frac{a}{b} \lambda^N$ for some $N \in \mathbb{N}$, the solution degenerates to a polynomial

$$f(z) = Q_\lambda \left(\frac{a}{b} \right) \sum_{n=0}^{N-1} \frac{(bz)^n}{n! Q_\lambda \left(\frac{a}{b} \lambda^n \right)},$$

In order to derive an expression for $f(z)$, which allows for determining its asymptotic behaviour, we recall the well known power series expansion

$$(4.3) \quad \frac{1}{Q_\lambda(\alpha)} = \sum_{n=0}^{\infty} (-1)^n \prod_{k=1}^n \frac{1}{1 - \lambda^k} \alpha^n$$

valid for $|\alpha| < 1$.

We now choose N as the smallest non-negative integer such that $|\frac{a}{b} \lambda^N| < 1$. Then we rewrite (4.2) as

$$(4.4) \quad f(z) = Q_\lambda \left(\frac{a}{b} \right) \sum_{n=0}^{N-1} \frac{(bz)^n}{n! Q_\lambda \left(\frac{a}{b} \lambda^n \right)} + Q_\lambda \left(\frac{a}{b} \right) \sum_{n=N}^{\infty} \frac{(bz)^n}{n! Q_\lambda \left(\frac{a}{b} \lambda^n \right)}.$$

We replace $\frac{1}{Q_\lambda}$ in the second sum by (4.3) to obtain

$$(4.5) \quad f(z) = Q_\lambda \left(\frac{a}{b} \right) \sum_{n=0}^{N-1} \frac{(bz)^n}{n! Q_\lambda \left(\frac{a}{b} \lambda^n \right)} + Q_\lambda \left(\frac{a}{b} \right) \sum_{m=0}^{\infty} (-1)^m \prod_{k=1}^m \frac{1}{1 - \lambda^k} \left(\frac{a}{b} \right)^m \sum_{n=N}^{\infty} \frac{(b \lambda^m z)^n}{n!},$$

which simplifies to

$$(4.6) \quad f(z) = Q_\lambda \left(\frac{a}{b} \right) \sum_{n=0}^{N-1} \frac{(bz)^n}{n! Q_\lambda \left(\frac{a}{b} \lambda^n \right)} + Q_\lambda \left(\frac{a}{b} \right) \sum_{m=0}^{\infty} (-1)^m \prod_{k=1}^m \frac{1}{1 - \lambda^k} \left(\frac{a}{b} \right)^m \left(e^{b \lambda^m z} - \sum_{n=0}^{N-1} \frac{(b \lambda^m z)^n}{n!} \right).$$

Remark 4.1. Notice that the expansion (4.6) converges for all values of a and b (after a suitable choice of N). This is in contrast to a similar series expansion given in [16], which converges only for $|a| < |b|$.

Remark 4.2. In [16] a generalization of the classical pantograph equation, the multi-pantograph equation

$$(4.7) \quad f'(z) = \sum_{k=1}^{\ell} a_k f(\lambda_k z) + b f(z)$$

with $1 > \lambda_{\ell} > \lambda_{\ell-1} > \cdots > \lambda_1 > 0$, $a_1, \dots, a_{\ell}, b \in \mathbb{R}$ is studied. In [16, Lemma 3.1] a series expansion for the

solution of (4.7) is given, which is shown to converge for

$$\sum_{k=1}^{\ell} |a_k| < |b|.$$

The truncation method presented above can easily be adapted to provide a series representation similar to (4.6) for f , which does not require this condition.

Acknowledgements. We are grateful to Daniel Berend for stimulating discussions and helpful remarks. G.D. thankfully acknowledge hospitality and support during his research visit in Graz, in May - June 2016, provided by Institut für Analysis und Zahlentheorie, Technische Universität Graz

REFERENCES

- [1] L. V. Bogachev, G. Derfel, and S. A. Molchanov, *On bounded continuous solutions of the archetypal equation with rescaling*, Proc. Royal Soc A471, 2015, 1–19.
- [2] N. G. de Bruijn, *The difference-differential equation $F'(x) = e^{ax+\beta} F(x-1)$. I, II*, Nederl. Akad. Wetensch. Proc. Ser. A. **56** = Indagationes Math. **15** (1953), 449–458, 459–464.
- [3] S. Chandrasekhar and G. Münch, *The theory of the fluctuations in brightness of the Milky Way, I*, *Astrophys. J.* **112**, (1950) 380–392.
- [4] G. Derfel, *On the asymptotics of the solutions of a class of functional-differential equations*, In: *Asymptotic Behavior of the Solutions of Functional- Differential Equations* (ed. A.N. Sharkovsky) , 1978, 58–66, Institute of Math. Ukrainian Acad. Sci Press, Kiev (in Russian).
- [5] G. Derfel, *Functional-differential equations with compressed arguments and polynomial coefficients: asymptotics of the solutions*, *J. Math. Anal. Appl.* **193**, (1995), 671–679.
- [6] G. Derfel, *Functional-differential and functional equations with rescaling*, *Operator Theory: Advances and Applications*, **80**, (1995), 100–111.
- [7] G. Derfel, P. J. Grabner, and F. Vogl, *Laplace operators on fractals and related functional equations (topical review article)*, *J. Phys. A.*, **46**, (2012), 463001-463034.
- [8] G. Derfel and A. Iserles, *The pantograph equation in the complex domain*, *J. Math. Anal. Appl.* **213**, (1997), 117–132.
- [9] T. S. Ferguson, *Lose a dollar or double your fortune*. In: *Proceedings of the 6th Berkeley symposium on mathematical statistics and probability, vol. III* (eds. L. M. Le Cam *et al.*), 1972, 657–666. Berkeley, CA: University of California Press.
- [10] D. Gaver Jr., *An absorption probability problem*. *J. Math. Anal. Appl.* **9**, (1964) 384–393.
- [11] P. J. Grabner and B. Steinsky, *Asymptotic behaviour of the poles of a special generating function for acyclic digraphs*, *Aequationes Math.* **70** (2005), 268–278.
- [12] A. Hall and G. Wake, *A functional differential equation arising in the modeling of cell growth*. *J. Austral. Math. Soc. Ser. B* **30**, (1989) 424–435.

- [13] A. Iserles, On the generalized pantograph functional-differential equation. *European J. Appl. Math.* (1993), 1-38
- [14] T. Kato and J. B. McLeod, *The functional-differential equation $y'(x) = ay(\lambda x) + by(x)$* , Bull. Amer. Math. Soc. **77** (1971), 891-937.
- [15] J. K. Langley, *A certain functional-differential equation*, J. Math. Anal. Appl. **244** (2000), no. 2, 564-567.
- [16] M. Z. Liu and D. Li, *Properties of analytic solution and numerical solution of multi-pantograph equation*, Appl. Math. Comput. **155** (2004), no. 3, 853-871.
- [17] W. Magnus, F. Oberhettinger, and R.P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer, Berlin, New York, 1966.
- [18] K. Mahler, On a special functional equation. *J. London Math. Soc.* **15**, (1940) 115-123.
- [19] G. R. Morris, A. Feldstein, and E. W. Bowen, The Phragmen - Lindelöf principle and a class of functional - differential equations, In: *Ordinary Differential Equations*, 1972, 513-540, Academic Press, San Diego.
- [20] J. R. Ockendon and A. B. Tayler, The dynamics of a current collection system for an electric locomotive. *Proc. Royal Soc. London A*, **322** (1971), 447-468.
- [21] R. W. Robinson, Counting labeled acyclic digraphs, In: *New Directions in the Theory of Graphs* (ed., F. Harari), 1973, 239-279 Academic Press, New York.
- [22] A. D. Sokal, The leading root of the partial theta function, *Advances in Mathematics*, **229**, no. 5, (2012), 2603-2621.
- [23] V. Spiridonov, Universal superpositions of coherent states and self-similar potentials. *Phys. Rev. A* **52**, (1995) 1909-1935.
- [24] C. Zhang, *On the solution to a certain functional differential equation*, J. Math. Anal. Appl. (2016), to appear, arXiv:1501.02700.

(G. D.) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, BEN GURION UNIVERSITY OF THE NEGEV, BEER SHEVA 84105, ISRAEL

E-mail address: derfel@math.bgu.ac.il

(P. G. & R. T.) INSTITUT FÜR ANALYSIS UND ZAHLENTHEORIE, TECHNISCHE UNIVERSITÄT GRAZ, KOPERNIKUSGASSE 24, 8010 GRAZ, AUSTRIA

E-mail address: peter.grabner@tugraz.at

E-mail address: tichy@tugraz.at